



**HAL**  
open science

## Perennial crops for sustainable soil management – Symbiotic fungi benefit from cultivation of a model crop for perennial cereal grain production in Europe.

Linda-Maria Dimitrova Mårtensson, Ana Barreiro, Olivier Duchene, Erik  
Steen Jensen

### ► To cite this version:

Linda-Maria Dimitrova Mårtensson, Ana Barreiro, Olivier Duchene, Erik Steen Jensen. Perennial crops for sustainable soil management – Symbiotic fungi benefit from cultivation of a model crop for perennial cereal grain production in Europe.. GLOBAL SYMPOSIUM ON SOIL BIODIVERSITY - FAO, 2020, Rome, Italy. hal-03653259

**HAL Id: hal-03653259**

**<https://isara.hal.science/hal-03653259>**

Submitted on 28 Apr 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Abstract template

---

GLOBAL SYMPOSIUM ON SOIL BIODIVERSITY | FAO HQ | Rome, Italy, 10-12 March 2020

## Perennial crops for sustainable soil management – Symbiotic fungi benefit from cultivation of a model crop for perennial cereal grain production in Europe

Linda-Maria Dimitrova Mårtensson<sup>1,\*</sup>, Ana Barreiro<sup>2</sup>, Olivier Duchene<sup>3</sup>, Erik Steen Jensen<sup>4</sup>

<sup>1</sup>Swedish University of Agricultural Sciences, Department of Biosystems and Technology, [Linda.Maria.Martensson@slu.se](mailto:Linda.Maria.Martensson@slu.se), +4640415129

<sup>2</sup>Swedish University of Agricultural Sciences, Department of Biosystems and Technology, [Ana.Barreiro@slu.se](mailto:Ana.Barreiro@slu.se)

<sup>3</sup>ISARA-Lyon, [olduchene@isara.fr](mailto:olduchene@isara.fr)

<sup>4</sup>Swedish University of Agricultural Sciences, Department of Biosystems and Technology, [Erik.Steen.Jensen@slu.se](mailto:Erik.Steen.Jensen@slu.se)

---

### Abstract summary

*Thinopyrum intermedium* (intermediate wheatgrass) is used as a model for future perennial cereal grain production and the influence of agricultural perennialization was assessed in terms of the abundance of arbuscular mycorrhizal fungi (AMF) in soil. *T. intermedium* proved to promote higher abundance of AMF in comparison to conventional, but also organic, annual wheat production as well as in comparison to annual wheat and rye under no-tillage cultivation. Our results indicate improved conditions for mycorrhizal fungi in perennial cereal production stands. We suggest that there may be a potential of this crop-microbe interaction to enhance nutrient status of the crop as well as crop performance, due to hyphal networks and improved soil structure.

*Keywords: Mycorrhizal fungi, soil fungi, soil ecology, perennial cereal grain crops, agriculture, soil ecosystem services*

---

### Introduction, scope and main objectives


Intensive agricultural practices are important drivers to the global loss of soil biodiversity and soil functions (Tsiafouli et al 2015). Thus, innovative management practices to improve soil biodiversity needs to be implemented in contemporary agriculture. *Thinopyrum intermedium* (intermediate wheatgrass) provide a perennial, multifunctional and multipurpose cereal crop (Duchene et al 2019). A shift from systems based on annuals to perennial grain-crop production with permanent to semi-permanent plant cover, without the negative consequences of inverting soil tillage, is expected to benefit the delivery of soil ecosystem services (Crews et al 2016). The soil microbial community structure and biomass can be used as early indicators of the effects of changed cropping practices on the soil quality. In particular, the hyphal growth of fungal organism groups, i.e. saprotrophic (Barreiro et al 2018) and mycorrhizal fungi (Hydbom et al 2017), benefit from less disturbance, and extensive management in general (Barreiro et al 2019). This would fulfil part of the second global Sustainable Development Goals (SDGs), calling for more resilient agricultural practices, enhancing both productivity and ecosystem quality. Increased soil biodiversity would also contribute to SDGs 12, 13 and 14 through improved efficiency in the management of natural resources, improved climate resilience as well as reduced nutrient loss. We determined the effect of agricultural perennialization of arable cropping systems by *T. intermedium* on the soil microbiome. We hypothesize that biomass of arbuscular mycorrhizal fungi (AMF) will be higher in the soil under perennial cereal crops compared to under annual cereals.

### Methodology

Mycorrhizal biomass indicators were analyzed in two field experiments. One field experiment was carried out at Alnarp (55.65°N, 13.06°E) in southernmost Sweden (established 2016), and another one in Maubec (50.37°N, 4.48°E) in southeast France (established in 2017). The Swedish experiment included sole crop stands of *T. intermedium* under organic management, intercrops of *T. intermedium* and *Medicago sativa*

(lucerne) under organic management, and *Tritium aestivum* (winter wheat) under conventional and organic management. The French experiment included conventional sole crop stands of *T. intermedium* in an annual grain crops rotation with *T. aestivum* (2017-2018), with direct sowing applied since 2007. Both experiments were replicated in four blocks for each treatment/crop. Soil sampling took place at the end of the growing season in 2018 using soil cores were four sub-samples made up one composite sample for each sampling depth and treatment plot.

Phospholipid fatty acid (PLFA) and neutral lipid fatty acid (NLFA) 16:1 $\omega$ 5 were used as indicators of AMF biomass (Olsson 1999; Frostegård and Bååth 1996; Olsson et al 1995). PLFA 16:1 $\omega$ 5 indicate both arbuscular mycorrhizal fungi (AMF) and Gram negative (G-) bacteria (Olsson 1999; Frostegård and Bååth 1996; Olsson et al 1995), thus, the NLFA 16:1 $\omega$ 5 is used as an accompanying indicator of AMF due to its high correlation with AMF spore counts in soil (Sharma and Buyer 2015). PLFAs and NLFAs were extracted using the protocol presented by Frostegård et al (1993) using CHCl<sub>3</sub>:MeOH:citrate buffer (1:2:0.8 v/v/v). The lipids were separated using pre-packed silica columns subsequently rinsed with CHCl<sub>3</sub> (neutral lipids) and MeOH (phospholipids). The neutral and phospholipids were subjected to methanolysis after adding FAME 19:0 as internal standard, and the resulting fatty acid methyl esters quantified by gas chromatography with flame ionization detector (GC-17A, Shimadzu).



**Figure 1:** I) The abundance of neutral lipid and phospholipid fatty acid indicators (nmol g<sup>-1</sup> dw soil) of arbuscular mycorrhizal fungi (AMF\_P and AMF\_N) in the conventional annual wheat (CON), organic annual wheat (ORG), organic perennial wheatgrass (PER), organic perennial wheatgrass and lucerne intercrop (PER IC) in the SITES AGROECOLOGICAL FIELD EXPERIMENT (SAFE) at Lönnstorp, SLU, Sweden. II) The abundance of AMF\_P and AMF\_N (nmol g<sup>-1</sup> dw soil) in perennial wheatgrass (PER) and annual wheat (ANN) at Maubec, France. Data are presented as means from each sampling depth (cm) and crop treatment with standard deviations. Different letters indicate differences at the significance level of  $p < 0.05$ . Upper case letters are used for AMF\_P and lower case letters for AMF\_N.

## Results

In the Swedish experiment, the abundance of the one of the biomass indicators of AMF was higher in the perennial sole crop than in the three other crops at the 0-30 cm soil level ( $F_{AMF\_N} = 13.48$  (df=3)  $p < 0.001$ ) (Fig. 1:I). The biomass indicators of AMF ( $F_{AMF\_P} = 112.6$  (df=1)  $p < 0.001$ ;  $F_{AMF\_N} = 38.54$  (df=1)  $p < 0.001$ ) decreased from 0-30 cm to 30-60 cm depth. In the French experiment, the abundance of PLFA biomass indicator of AMF was higher in the perennial sole crop than in the annual crops at the level of 0-10 and 10-

30 cm soil depth ( $F_{AMF,P}=4.441$  (df=1)  $p<0.05$ ) (Fig. 1:II). The biomass indicators of AMF was generally higher in the upper soil level than further down the soil profile ( $F_{AMF,P}=121.1$  (df=3)  $p<0.001$ ;  $F_{AMF,N}=6.516$  (df=3)  $p<0.01$ ).

## Discussion

Our hypothesis was partly confirmed in the Swedish experiment through the higher abundance of the NLFA marker for AMF biomass. The lack of influence on the PLFA marker for AMF biomass may be due to the rather young age of the perennial stand. The high abundance, and clear response, of the NLFA AMF marker reflect the larger abundance of spores AMF spore counts (Sharma and Buyer 2015) in soil under the perennial crop. The slightly higher abundance of the NLFA marker of AMF biomass in organic wheat may be an early result of zero use of fungicides. The French experiment confirms the expectation to increase abundance of AMF biomass in soil under the perennial crop. It has been shown that AMF colonization increases with increasing age of co-occurring plant (Grigera et al. 2007, Zimmermann et al. 2016). Further questions to explore is the potential to increase the plant's ability to acquire nutrients (Tran et al 2019) and the potential of AMF to improve soil structure and related soil functions. Intercropping with legumes, to exploit biological  $N_2$  fixation, is an important measure to contribute to sustainable development (Voisin et al, 2014) through increased yield stability (Raseduzzaman & Jensen 2017) and the potential higher biomass N concentration and N accumulation in the intercropped partner (Boller and Nösberger, 1987). The occurrence of  $N_2$  fixing bacteria and the higher levels of accessible soil N usually benefit the abundance of AMF. Our results indicate that the expected higher abundance of AMF under perennial crops is not realized intercropped with lucerne, which could be explained the competitive interactions resulting in less *T. intermedium* roots for the AMF to colonize. Furthermore, the otherwise expected AMF colonization of lucerne roots may not have occurred and the interactions needs to be further investigated.

## Conclusions

Agricultural perennialization will lead to healthier soil microbiomes, in this case proven by higher abundance of the obligate symbiotic soil mycorrhizal fungi. The outcome is in line with the global sustainability goals meaning that the implementation of improved, less interruptive agricultural practices should be promoted.

## Acknowledgements

---

Part of this study was carried out at the SITES Lönnstorp Research Station, at the Swedish University of Agricultural Sciences (SLU), Alnarp, Sweden.

---

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

## References

- Barreiro, A., Fox, A., Widmer, F., Lüscher, A., Zimmermann, J., Musyoki, M., Rasche, F., Dimitrova Mårtensson, L-M.** 2018. Soil microbial communities in highland and lowland grasslands under different management intensity. Focused meeting of the Microbiology Society 'Microbiomes underpinning agriculture'. Cork, Ireland. Conference abstract.
- Barreiro, A., Fox, A., Lüscher, A., Widmer, F., Vieira, A., Parelho, C., Silva, L., Melo, J., Cruz, C., Musyoki, M., Zimmermann, J., Rasche, F., Dimitrova Mårtensson, L-M.** 2019. Soil microbial biomass and community structure in grasslands along a European gradient In: Improving sown grasslands through breeding and management, Vol 24 Grassland Science in Europe, pp 464.
- Boller, B.C. & Nösberger, J.** 1987. Symbiotically fixed nitrogen from field- grown white and red clover mixed with ryegrasses at low levels of 15-N-fertilization. Plant and Soil 104:219-226.

- Crews, T.E., Blesh, J., Culman, S.W., Hayes, R.C., Jensen, E.S., Mack, M.C., Peoples, M.B. & Schipanski, M.E. 2016. Going where no grains have gone before: From early to mid-succession. *Agriculture Ecosystems and Environment* 223:223-238.
- Duchene, O., Celette, F., Ryan, M. R., DeHaan, L.R., Crews, T.E, David, C. (2019) Integrating multipurpose perennial grains crops in Western European farming systems. *Agriculture, Ecosystems and Environment* 284.
- Frostegård, Å., Bååth, E., & Tunlid, A. 1993. Shifts in the structure of soil microbial communities in limed forests as revealed by phospholipid fatty acid analysis. *Soil Biology and Biochemistry* 25:723-730.
- Frostegård, Å. & Bååth, E. 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. *Biology and Fertility of Soils* 22:59-65.
- Hydbom, S., Ernfors, M., Birgander, J., Hollander, J., Jensen, E.S., & Olsson, P.A. 2017. Reduced tillage stimulated symbiotic fungi and microbial saprotrophs, but did not lead to a shift in the saprotrophic microorganism community structure. *Applied Soil Ecology* 119:104-114.
- Grigera, M., Drijber, R. A., Wienhold, B. J. 2007. Increased abundance of arbuscular mycorrhizal fungi in soil coincides with the reproductive stages of maize. *Soil Biology and Biochemistry* 39:1401-1409.
- Zimmermann, J., Musyoki, M. K., Cadisch, G., Rasche, F. 2016. Biocontrol agent *Fusarium oxysporum* f.sp. *strigae* has no adverse effect on indigenous total fungal communities and specific AMF taxa in contrasting maize rhizospheres. *Fungal Ecology* 23:1-10.
- Kladivko, E.J. 2001. Tillage systems and soil ecology. *Soil Tillage Research* 61:61-76.
- Olsson, P.A. 1999. Signature fatty acids provide tools for determination of the distribution and interactions of mycorrhizal fungi in soil. *FEMS Microbiology Ecology* 29:303-310.
- Olsson, P.A., Bååth, E., Jakobsen, I., & Söderström, B. 1995. The use of phospholipid and neutral lipid fatty acids to estimate biomass of arbuscular mycorrhizal fungi in soil. *Mycology Research* 99:623-629.
- Raseduzzaman, Md. & Jensen, E.S. 2017. Does intercropping enhance yield stability in arable crop production? A meta-analysis. *European Journal of Agronomy* 91:25-33.
- Sharma, M.P. & Buyer, J.S. 2015. Comparison of biochemical and microscopic methods for quantification of arbuscular mycorrhizal fungi in soil and roots. *Applied Soil Ecology* 95, 86-89.
- Tsiafouli, M.A., Thébault, E., Sgardelis, S.P., de, Ruiter, P.C., van der, Putten, W.H., Birkhofer, K., Hemerik, L., de, Vries, F.T., Bardgett, R.D., Brady, M.V., Bjornlund, L., Jørgensen, H.B., Christensen, S., Hertefeldt, T.D., Hotes, S., Gera Hol, W., Frouz, J., Liiri, M., Mortimer, S.R., Setälä, H., Tzanopoulos, J., Uteseny, K., Pižl, V., Stary, J., Wolters, V. & Hedlund, K. 2015. Intensive agriculture reduces soil biodiversity across Europe. *Global Change Biology* 21:973-985.
- Tran, B.T.T., Watts-Williams, S.J., & Cavagnaro, T.R. 2019. Impact of an arbuscular mycorrhizal fungus on the growth and nutrition of fifteen crop and pasture plant species. *Functional Plant Biology* 46:732-742.
- Voisin, A-S., Guéguen, J., Huyghe, C., Jeuffroy, M-H., Magrini, M-B., Meynard, J-M, Mougél, C., Pellerin, S. & Pelzer, E. 2014. Legumes for feed, food, biomaterials and bioenergy in Europe: a review. *Agronomy for Sustainable Development* 34:361-380.